

## Waste land management with water chestnut a policy development for policy planning and research in Hardoi, U.P., India

Trilok Nath Rai<sup>1</sup>, Kedar Nath Rai<sup>2</sup>, Sanjeev Kumar Rai<sup>3</sup>, Rajeev Kumar Rai<sup>4</sup>, Mohit Singh<sup>5</sup>

<sup>1</sup> Department of Soil Science & Agricultural Chemistry, Krishi Vigyan Kendra (ICAR-CSSRI), Dhikunni, Sandila, Hardoi, Uttar Pradesh, India

<sup>2</sup> Associate Professor and Head, Department of Soil Science & Agricultural Chemistry, Kamla Nehru Institute of Physical and Social Sciences, Sultanpur, Uttar Pradesh, India

<sup>3</sup> Department of Soil Science & Agricultural Chemistry, Kamla Nehru Institute of Physical and Social Sciences, Sultanpur, Uttar Pradesh, India

<sup>4</sup> Department of Nursing, Institute of Nursing & Paramedical Sciences Sultanpur, Uttar Pradesh, India

<sup>5</sup> Agriculture Extension, Krishi Vigyan Kendra (ICAR-CSSRI), Dhikunni, Sandila, Hardoi, Uttar Pradesh, India

### Abstract

Water chestnut is an important high value aquatic crop grown in India. Of late the importance of this crop has been increasing due to the presence of appreciable amount of nutritional contents therein like protein, carbohydrates, fiber and minerals. This paper explores the recent agronomic management techniques and post-harvest utilization of water chestnut crop in Saline – Sodic Soils of Hardoi, U.P. Being a short duration annual crop, it can be introduced in low-input cropping systems as a viable alternative crop for sustainable horticulture. Water chestnut production is mainly constrained by lack of knowledge on agronomic practices, deficiency of technical know-how about its processing and lack of mass awareness about its high nutritional values. These days, field cultivation of this crop is also proving to be profitable. Hence, attempts are being undertaken to integrate it with field crops. Improved genotypes namely Bihar Large Red, Green Spineless and Balasore Green have been found to perform the best under field cultivation. Planting distance of 1m x 2m with application of NPK @ 60:40:30 kg/ha in split applications have witnessed a yield of as much as 12.0 t/ha. Being a high value cash crop, it can be sold as the fresh produce; better income however can be secured by selling value added products of water chestnut like dry nut and flour. Cultivated with right agronomic techniques and processed appropriately after harvest, water chestnut crop can significantly improve the livelihood of farmers living in water surplus regions of rural India.

**Keywords:** Saline–sodic soils, bihar large red, green spineless, balasore green, hardoi

### Introduction

Water chestnut (*Trapa natans* L.) is one of the most important aquatic nuts grown in India. It is an important nut in terms of protein, carbohydrates, mineral and vitamin content along with important amino acids (Faruk *et al.*, 2012) [6]. Nutritional and medicinal qualities of *Trapa natans* have been recognized in India, China, Thailand, Russia and many other countries (Hummel and Kiviat, 2004) [8].

In India, *Trapa natans*, *T. bispinosa*, *T. quadrispinosa* and *T. bicornis* are very popular. Fresh nut is well known for its high water content (80.0%) (Puste, 2004) [14] protein (1.87%) and TSS (7.0-8.0%) (Singh *et al.*, 2010) [17]. According to Adkar *et al.*, (2014) [1] and Alfasane *et al.*, (2011) [2] protein and carbohydrate content of fresh water chestnut are 4.40% and 22.3%, respectively. Water chestnut is an excellent source of crude fiber (2.13% in green variety and 2.27 % in red variety). It is also a fair source of Vitamin B along with Ca, K, Fe and Zn (Singh *et al.*, 2009 and Gopalan *et al.*, 2004) [7, 16]. Water chestnut is an eco-friendly crop and is utilized for prevention of cyano-bacterial bloom and addition of nutrients to the soil after decomposition of its residue. *Trapa natans* usually decreases dissolved inorganic nitrogen in water thereby potentially reducing eutrophication process in water bodies (Tsuchiya and Iwakuma, 1993) [18]. From aquatic environment, it absorbs heavy metals such as cadmium, nickel, and cobalt (USEPA, 1989) [19] which are deposited mainly in hulls. According to Block and Morris (2002) [4] one seed gives rise to 10-12

rosettes and each rosette produces as much as 20 fruits. It requires 4.0 months after transplanting for producing economic yield. It is very hardy crop requiring only 1-2 sprays of insecticides and fungicides for disease and pest free plant growth. This crop cultivation is maximum in salt affected area of Hardoi, Uttar-Pradesh.

### Agronomic management techniques Genotype

Four types of water chestnut are mainly cultivated in India *viz.* spine and spineless with red and green color. The prominent genotypes of water chestnut are Spineless Red, Spineless Green (Jana *et al.*, 2019) [10, 11]. Red Spine, Bihar Large Red, Green Spine, Balia Red, Lucknow Green, Jaunpuri and Balasore Green (Jana, 2016a) [9] which are common in all parts of India. However Balasore Green of Odisha gives relatively a better yield of 16.84 t/ha (Choudhary *et al.*, 2003) [5].

### Climate

Water chestnut needs rather cool climate for its seed germination (12-15 °C). For plant growth and development it needs little higher temperature 25-30°C. However, fruit is generally harvested during cool winter October-November (Block and Morris, 2002 and Kundu and Joshi, 2012) [4, 13].

### Soil and nutrient management

It thrives in soft nutrient rich waters of lakes, ponds and streams with a neutral to slightly alkaline pH. For commercial cultivation soil should be heavy, muddy and

rich in organic matter. Organic manures in form of oil cakes, poultry manure and compost/FYM @ 5-8 t/ha are recommended for growth and development of water chestnut crop (Choudhary *et al.*, 2003) [5]. Significant amount of phosphorus and potassium is needed for optimum growth of the crop. In Uttar-Pradesh, application of 35-45 kg urea/ha during transplantation and another 25 kg/ha is recommended after 25-35 days of transplanting (Jana, 2016a) [9]. Soils rich in Zn and B result in increased number of fruits and fruit weight. For newly cultivated field, the fertilizer dose can be modified from NPKS @ 65:25:25:25 kg/ha to NPKS @ 65:45:35:25 kg/ha (Jana *et al.*, 2019) [10, 11].

### Nursery raising and transplanting

The seed rate for nursery preparation is 80-100 kg dry nut/ha. It is recommended that plants are first grown in a low nutrient plot/nursery and transplanted into main plot when stems are about 1.0-1.5 m tall. Care should be taken at the time of transplanting to keep the seedlings moist but not submerged. According to Jana *et al.*, (2019) [10, 11] planting distance should be maintained at 1m x 2m to give an optimum plant population of 4500- 5000 plants/ha.

### Water level

Throughout the growth period of the crop 1-2 m water depth should be maintained for proper growth and development (Jana, 2016a) [9]. A greater depth of water is tolerated by the plants but they do not prosper if water shortage occurs at early period of growth. Prior to harvesting of the crop water depth may be reduced gradually.

### Reproductive stage

Spineless Red cultivar flowers first at wetland ecosystem of Uttar Pradesh during 3<sup>rd</sup> week of August. Blooming period continues to 2<sup>nd</sup> week of September. But Spineless Green cultivar has longer blooming period from August to October. Harvesting of the nuts is usually performed during October to December. A well maintained local genotype gives the maximum yield of 2.75-3.95 t/ha (Jana, 2016) [9]. Bihar Large Red (Spine), the commercial variety of Mithila region, Darbhanga, India, gives a higher yield of 7.0 t/ha (Jana *et al.*, 2019) [10, 11]. The maximum yield of 12.50 t/ha can be achieved by cultivation of Green Spineless genotype (Jana *et al.*, 2019) [10, 11]. Regular eradication of aquatic weeds, especially *Hydrilla* and *Sagittaria*, common in water chestnut crop, is utmost important during the cropping season to avoid the crop-weed competition (Jana, 2016b) [9].

### Pest and disease management

Beetle (*Galerucella birmanica* Jacoby) and aphid are the main pests under wetland ecosystem of Uttar Pradesh. Two to three sprays of *Carbaryl* @ 0.075% at an interval of 5-10 days can control beetle (*Galerucella birmanica* Jacoby), aphid and mealy bug. Spray of *Bavistin* (0.15%) is required to control fungal diseases (Yadav and Gargav, 1988) [20].

### Value addition

Traditionally water chestnut is consumed after boiling. This is high value nut from which sweets and precious chapatti may be made. Sun drying or artificial drying in oven is

common in India. It's a medium to low calorie food with maximum nutrients which is gaining importance for making snacks for every class of people throughout the world. Nut is free from gluten and cholesterol and rich in K, Mg and fiber, which renders it an ideal snack. Low cost mixed food can be made by mixing this nut flour with wheat and pulse flour to fortify its nutritional value (Sha *et al.*, 2016; and Baba *et al.*, 2016) [3, 15].

### Water chestnut for enhancing rural livelihood sustainability

Water chestnut is a high value aquatic horticultural cash crop. High farmers' income can be secured by increasing production of water chestnut by cultivating high yielding genotypes, value addition and efficient marketing management. The net profit from water chestnut (fresh product) is about INR

2.75 lakh/ha/year (Jana, 2019) [10, 11] with production cost of about INR 0.55 lakh/ha/year. From the cultivation of superior genotype 'Green Spineless', a harvest of more than 2.5 t/ha dry nut is possible which is worth INR 4.55 lakh/ha/year.

### Limitations and future prospects

Necessary actions should be undertaken to keep ecosystem vibrant and get the maximum benefit from water chestnut crop. Crop residues are easily decomposed and add various plant nutrients to soil including organic matter for sustainable horticultural production system. Water chestnut cultivation is also a kind of bio-remediation by natural means. It is difficult to maintain its quality during processing which is essential for export, which merits particular attention. Water chestnut is a lucrative crop which requires mass awareness about nutritional qualities and consumption techniques for the health conscious people. Water chestnut crop can significantly improve the livelihood of farmers living in water surplus regions of rural India if cultivated with right agronomic techniques and processed appropriately after harvest.

**Success story:** This is success story of Mr. Vimal Kumar (age 22 years) s/o Shri Bablu Kashyap, who lived in dhikunni, bharawan, U.P., India. Basically he is marginal farmers who depends on water chest nut and fish farming since last five years. He joined KVK, Hardoi-II during 2021-2022 with SMS, Soil Science Dr. T.N.Rai and taken guidance for cultivation of water chest nut.

**Conclusions:** This is concluded that this crop is suitable under saline - sodic soils of India which is found in maximum areas of India.

### Conflicts of interests

Authors have declared that they have no known competing financial interests or non-financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Acknowledgements:** Author Dr. T.N. Rai thankful to corresponding author Dr. K.N. Rai.



Water chestnut dried ( Mr. Vimal Kumar) for sale @ 12500.0- 15500.0 per q. at Dhikunni, Bharawan, Hardoi,U.P. during 2025-2026, survey and awareness works done by SMS, Soil Science from 2021-2026 at Hardoi.



### References

1. Adkar P, Dongare A, Ambavade S, Bhaskar VH. *Trapabis pinosa* Roxb.: A review on nutritional and pharmacological aspects, 2014.
2. Alfasane M, Almujaadade M, Khonder M, Rahman M. Biochemical composition of the fruits of water chestnut (*Trapabis pinosa* Roxb.). *Dhaka Univ. J. Biol. Sci.*,2011:20(1):95-98.
3. Baba WN, Din S, Punoo HA, Wani TA, Ahmad M, Masoodi FA. *et al.* Comparison of cheese and paneer whey for production of a functional pineapple beverage: nutraceutical properties and shelf life. *J. Food. Sci. Technol.*,2016:53:2558-2568.
4. Block AF, Morris TA. Arboretum of the University of Pennsylvania. European water-chestnut *Trapa natans* L. Water chestnut family (Trapaceae), 2002.
5. Choudhary SR, Sahoo N, Verma N. Growth behavior and yield of five water chestnut varieties under waterlogged condition. *Indian J. Plant Physiol*,2003:8(4):369-371.
6. Faruk MO, Amin MZ, Sana NK, Saha RK, Biswas RK. Biochemical analysis of two varieties of water chestnuts (*Trapa* spp). *Pakistan Journal of Biological Sciences*,2012:15(21):1019-1026.
7. Gopalan C, Sastri R, Balasubramanian SC. Nutritive value of Indian foods, 2004.

8. Hummel M, Kiviat E. Review of world literature on water chestnut with implications for management in North America. *J Aquat Plant Manage*,2004;42:17-27.
9. Jana BR. The impact of water depth on growth of gorgon nut (*Euryale ferox* Salisb.) and associated weeds under field cultivation. *Journal of Crop and Weed*,2016;13(1):77-82.
10. Jana BR. Exploring aquatic horticultural crops for higher income of farmers living in water surplus ecology, 2019, 61-70.
11. Jana BR, Bhatt BP, Singh IS, Idris M. A study on commercial cultivation and storage of water chestnut (*Trapa natans* L.) under wetland ecosystem of North Bihar, India. *Journal of Applied and Natural Science*,2019;11(2):528-533.
12. Jana H. Water Caltrop: A potential crop of water bodies. *Rastriya Krishi*,2016;11(2):7-11.
13. Kundu M, Joshi R. Production technology of water chestnut (*Trapa natans*), 2012, 1-2.
14. Puste AM. Agronomic management of wetland crops, 2004.
15. Sha M, Baba WN, Masoodi FA, Bazaz R. Wheat-water chestnut flour blends: effect of baking on antioxidant properties of cookies, 2016.
16. Singh GD, Singh ASB, Saxena DC. Physicochemical, pasting, thermal and morphological characteristics of Indian water chestnut (*Trapa natans*) starch. *Starch Stärke*,2009;61:35-42.
17. Singh GD, Singh S, Jindal N, Saxena DC. Physico-chemical characteristics and sensory quality of singhara (*Trapa natans* L.): An Indian water chestnut under commercial and industrial storage conditions. *Afr. J. Food Sci*,2010;4(11):693-702.
18. Tsuchiya T, Iwakuma T. Growth and leaf life-span of a floating-leaved plant, *Trapa natans* L., as influenced by nitrogen flux. *Aquatic Botany*,1993;46(3-4):317-324.
19. USEPA. Superfund record of decision, Marathon Battery, NY, third remedial action - final, 1989.
20. Yadav HS, Gargav VP. Chemical control of singhara beetle, *Galerucella Birmanica* Jacoby. *Indian Journal of Plant Protection*,1988;16:159-162.